

Naloxone for Overdose Prevention

patient name

date of birth

patient address

patient city, state, ZIP code



prescriber name

prescriber address

prescriber city, state, ZIP code

prescriber phone number

Naloxone HCl 0.4 mg/mL (Narcan)

1 x 10 mL as one fliptop vial (NDC 0409-1219-01) OR
2 x 1mL single dose vials (NDC 0409-1215-01)

Refills: _____

Intramuscular (IM) syringe, 23 G, 3cc, 1 inch

Qty: _____

Refills: _____

Sig: For suspected opioid overdose,
inject 1mL IM in shoulder or thigh.

Repeat after 3 minutes if no or minimal response.

prescriber signature

date

Detach for patient



Are they breathing?

Signs of an overdose:

- Slow or shallow breathing
- Gasping for air when sleeping or weird snoring
- Pale or bluish skin
- Slow heartbeat, low blood pressure
- Won't wake up or respond (rub knuckles on sternum)



Call 911 for help

All you have to say:

"Someone is unresponsive and not breathing."

Give clear address and location.



Airway

Make sure nothing is inside the person's mouth.



Rescue breathing

Oxygen saves lives. Breathe for them.

One hand on chin, tilt head back, pinch nose closed.

Make a seal over mouth & breathe in

1 breath every 5 seconds

Chest should rise, not stomach



Evaluate

Are they any better? Can you get naloxone
and prepare it quickly enough that they won't
go for too long without your breathing assistance?



Prepare naloxone

- Remove cap from naloxone and uncover needle
- Insert needle through rubber plug, with bottle upside down
- Pull back on plunger and take up 1 cc into the syringe
- Don't worry about air bubbles (they aren't dangerous in muscle injections)



Muscular injection

inject 1cc of naloxone into a big muscle (shoulder or thigh)



Evaluate + support

- Continue rescue breathing
- Give another shot of naloxone in 3 minutes if no or minimal breathing or responsiveness
- Naloxone wears off in 30-90 minutes
- Comfort them; withdrawal can be unpleasant
- Get them medical care and help them not use more opiate right away
- Encourage survivors to seek treatment if they feel they have a problem

How to Avoid Overdose

- Only take medicine prescribed to you
- Don't take more than instructed
 - Call a doctor if your pain gets worse
- Never mix pain meds with alcohol
- Avoid sleeping pills when taking pain meds
- Dispose of unused medications
- Store your medicine in a secure place
- Learn how to use naloxone
- Teach your family + friends how to respond to an overdose

For More Info
PrescribeToPrevent.com

Poison Center
1-800-222-1222
(free & anonymous)